

## Package insert

# Traumeel ad us. vet. gel

## 1. Scheduling status

Schedule 0

## 2. Proprietary name and dosage form

Traumeel ad us. vet. gel for topical use.

## 3. Composition

100g cont.: Achillea millefolium D0 0,09g; Aconitum napellus D1 0,05g; Arnica montana D3 1,5 g; Atropa bella-donna D1 0,05g; Bellis perennis D0 0,1g; Calendula officinalis D0 0,45g; Echinacea D0 0,15g; Echinacea purpurea D0 0,15g; Hamamelis virginiana D0 0,45g; Hypericum perforatum D6 0,09g; Matricaria recutita D0 0,15g; Symphytum officinale D4 0,1g; Hepar sulfuris D6 0,025g; Mercurius solubilis Hahnemanni D6 0,04g.  
Excipients: Purified water, Ethanol (96 percent).

## 4. Pharmacological classification

D. 33.2. Homeopathic preparations

Discipline of the medicine: Homeopathy

## 5. Pharmacological action

Traumeel ad us. vet. is prepared in accordance with homeopathic principles.

## 6. Target species

Cattle, horse, donkey, pig, sheep, goat, dog, cat and small pets.

## 7. Indications

This medicine is proposed to treat lameness and pain caused by blunt injuries such as sprains, contusions and bruises, and traumatic or surgical wounds.

Use in the treatment of inflammatory diseases of the musculoskeletal system (e.g. arthritis, bursitis, tenosynovitis) as well as inflammatory diseases of other tissues (e.g. mastitis, dermatitis, phlegmon).

## 8. Contraindications

Do not use in animals with known hypersensitivity to one of the ingredients of the veterinary medicinal product.

## 9. Special warnings

### Special warnings for each target species

For animal treatment only. Medicinal products must be kept out of the reach of children. Do not administer near or into eyes, on mucous membranes or injured skin. This product contains ethanol.

### Special precautions for safe use in the target species

If symptoms in the animal persist or worsen, you should consult a veterinarian.

## 10. Interactions

As with all medicinal products, homeopathic veterinary medicinal products may interact with other veterinary medicinal products. Please inform the attending veterinarian if your animal receives any other medicines at the same time.

## 11. Pregnancy and lactation

The safety of the veterinary medicinal product has not been established during pregnancy, lactation or laying period. Like all medicines, this product should only be used during pregnancy or lactation after consultation with the veterinarian.

## 12. Dosage and directions for use

### Method of administration

For external use only.

Unless otherwise prescribed:

Apply the gel several times daily (up to 3–4 times) to the affected area and rub in gently until completely resorbed. Continue with this regime until symptoms have dissipated or are resolved. The product may be applied using mild compression bandaging and/or occlusive bandaging.

### **13. Side effects**

Homeopathic veterinary medicinal products are generally well tolerated. In rare cases, side effects may occur, although not every animal gets them.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. If you notice any side effect(s), listed or not, or you think that the medicine has not worked, please contact your veterinarian.

### **14. Known symptoms of over dosage and particulars of its treatment**

In case of accidental administration or ingestion of quantities exceeding the recommended dosage, no special countermeasures are required. A veterinarian may be consulted as a precaution in case of overdose.

### **15. Withdrawal periods**

Cattle, horses, donkeys, pigs, sheep, goats:

Meat: zero days

Cattle, horses, donkeys, sheep, goats:

Milk: zero hours

### **16. Identification**

Tube (aluminium) with 50 g gel and plastic bottle with 250 g gel.

### **17. Storage instructions**

Store at or below 30 °C, away from children and uninformed persons.

### **18. Registration number**

To be allocated by the SAHPRA.

### **19. Name and business address of the holder of the certificate of registration**

ModHomCo (Pty) Ltd  
96 Amsterdam Street, Clubview, 0157 Centurion  
Manufactured in Germany.

### **20. Date of publication of the package insert**

January 2026

This unregistered medicine has not been evaluated by the SAHPRA for its quality, safety or intended use.

## **Inligtingstuk**

# **Traumeel ad us. vet. gel**

### **1. Skeduleringstatus**

Skedule 0

### **2. Handelsnaam en farmaseutiese vorm**

Traumeel ad us. vet. gel vir plaaslike aanwending.

### **3. Samestelling**

100g bevat.: Achillea millefolium D0 0,09g; Aconitum napellus D1 0,05g; Arnica montana D3 1,5 g; Atropa bella-donna D1 0,05g; Bellis perennis D0 0,1g; Calendula officinalis D0 0,45g; Echinacea D0 0,15g; Echinacea purpurea D0 0,15g; Hamamelis virginiana D0 0,45g; Hypericum perforatum D6

0,09g; Matricaria recutita D0 0,15g; Symphytum officinale D4 0,1g; Hepar sulfuris D6 0,025g; Mercurius solubilis Hahnemanni D6 0,04g.  
Bymiddels: Gesuiwerde water, Etanol (96 persent).

#### **4. Farmakologiese klassifikasie**

D. 33.2. Homeopatiese preparate

Dissipline van die medisyne: Homeopatie

#### **5. Farmakologiese werking**

Traumeel ad us. vet. word volgens homeopatiese beginsels voorberei.

#### **6. Teikenspesies**

Beeste, perd, donkie, vark, skape, bok, hond, kat en klein troeteldiere.

#### **7. Indikasies**

Hierdie medisyne word voorgestel om mankheid en pyn te behandel wat veroorsaak word deur beserings soos verstuitings, kneusings en kneusplekke, en traumatiese of sjirurgiese wonde. Gebruik in die behandeling van inflammatoriese siektes van die muskuloskeletale stelsel (bv. artritis, bursitis, tenosinovitis) sowel as inflammatoriese siektes van ander weefsels (bv. mastitis, dermatitis, flegmon).

#### **8. Kontra-indikasies**

Moenie gebruik word by diere met bekende hipersensitiwiteit vir een van die bestanddele van die veeartsenykundige middel nie.

#### **9. Spesiale waarskuwings**

Spesiale waarskuwings vir elke teikenspesie

Slegs vir dierebehandeling. Medisyne moet buite bereik van kinders gehou word. Moenie naby of in oë, op slymvliese of beseerde vel toedien nie. Hierdie produk bevat etanol.

Spesiale voorsorgmaatreëls vir veilige gebruik in die teikenspesie

As simptome by die dier voortduur of vererger, moet u 'n veearts raadpleeg.

#### **10. Interaksies**

Soos met alle medisinale produkte, kan homeopatiese veeartsenykundige produkte interaksie met ander veeartsenykundige produkte hê. Stel asseblief die behandelende veearts in kennis indien u dier enige ander medisyne op dieselfde tyd gebruik.

#### **11. Swangerskap en laktasie**

Die veiligheid van die veeartsenykundige produk is nie vasgestel tydens dragtigheid, laktasie of lêperiode nie. Soos alle medisyne, moet hierdie produk slegs tydens dragtigheid of laktasie gebruik word na oorlegpleging met die veearts.

#### **12. Dosis en gebruiksaanwysings**

Metode van toediening:

Alleenlik vir uitwendige gebruik.

Tensy anders voorgeskryf:

Wend die gel 'n paar keer per dag (tot 3-4 keer) aan op die aangetaste area en vryf dit saggies in totdat dit heeltemal geabsorbeer is. Hou aan met die behandelingsregime totdat simptome verbeter of verdwyn het. Die produk kan met 'n ligte kompressieverband en/of okklusiewe verband aangewend word.

#### **13. Nuwe-effekte**

Homeopatiese veeartsenykundige medisyne word oor die algemeen goed verdra. In seldsame gevalle kan nuwe-effekte ervaar word, hoewel dit nie by alle diere voorkom nie.

Dit is belangrik om nadelige gebeurtenisse te rapporteer. Dit verseker deurlopende veiligheidsmonitering van die veeartsenykundige produk. As jy enige nuwe-effek(e) opmerk, hetsy dit in hierdie inligtingstuk gemeld is of nie, of jy dink dat die medisyne nie gewerk het nie, kontak asseblief jou veearts.

**14. Bekende simptome van oordosering en besonderhede van die behandeling daarvan**

In die geval van toevallige toediening of inname van hoeveelhede wat die aanbevole dosis oorskry, is geen spesiale teenmaatreëls nodig nie. 'n Veearts kan as 'n voorsorgmaatreël geraadpleeg word in geval van oordosis.

**15. Onttrekkingsperiodes**

Beeste, perde, donkies, varke, skape, bokke:

Vleis: nul dae

Beeste, perde, donkies, skape, bokke:

Melk: nul ure

**16. Identifikasie**

Buis (aluminium) met 50 g gel en plastiekbottel met 250 g gel.

**17. Opbergingsinstruksies**

Bêre by of onder 30 °C, weg van kinders en oningeligte persone.

**18. Registrasienommer**

Om deur die SAHPRA toegeken te word.

**19. Naam en besigheidsadres van die houer van die registrasiesertifikaat**

ModHomCo (Edms) Bpk

Amsterdamstraat 96, Clubview, 0157 Centurion

Vervaardig in Duitsland.

**20. Datum van publikasie van die voubiljet**

Januarie 2026

Hierdie ongeregistreerde medisyne is nie deur die SAHPRA vir die kwaliteit, veiligheid of beoogde gebruik geëvalueer nie.